

DOES SHORT SEGMENT LUMBAR STABILIZATION AND FUSION ACCELERATE ADJACENT UPPER SEGMENT INSTABILITY?

KISA SEGMENT LOMBER STABİLİZASYON VE FÜZYON KOMŞU SEGMENT İNSTABİLİTESİNİ HIZLANDIRIR MI?

Emin Kasım*, Uygur Er**, Serkan Şimşek***, Atilla Kazancı*,
Burak Kazancı****, Bülent Güçlü****, Murad Bavbek*

ABSTRACT:

Objective: The true incidence and clinical impact of instability at the adjacent segment after stabilization of the lumbar spine was unclear. This study investigates development of disc herniation or instability at the adjacent segment after stabilization of the lumbar spine.

Methods: Twenty-nine patients who were operated upon using short segment posterior instrumentation and fusion were evaluated for development of adjacent upper segment disc herniation and instability. Stabilization and fusion were performed for listhesis or the presumed instability after stenosis operation or unstable disc herniation. In all patients, pre- and postoperative MRIs and X-ray images were obtained. In MRI, disc herniations were evaluated. In X-ray, slippage and rotational angling of the adjacent upper segment were measured.

Results: Three patients had disc herniation and six patients had instability at the adjacent segment. No statistical correlation was found between development of instability and disc herniation and age, gender of the patient, type of the pathology, or the level stabilized.

Conclusions: Stabilization and fusion were sufficient for the patients and did not produce too much instability at the adjacent upper segment. Randomized studies with more patients and different instability criteria are needed to better understand adjacent segment instability.

Keywords: Adjacent segment instability, degenerative lumbar spine disease, Instrumented fusion, short segment fusion

Level of evidence: Retrospective clinical study, Level III

ÖZET

Amaç: Lomber omurga sabitlemesinden sonra gelişen komşu segment instabilitesinin gerçek insidansı ve kliniğe yansımaları aydınlatılmamıştır. Bu çalışma lomber omurga sabitlemesinden sonra komşu segmentte disk hernisi veya instabilite gelişiminin incelenmesi amaçlanmıştır.

Yöntem: Kısa segment posterior enstrümantasyon ve füzyon uygulanan 29 hasta komşu segmentte disk hernisi veya instabilite gelişimi açısından incelenmiştir. Sabitleme ve füzyon listezis, instabil disk hernisi veya stenoz ameliyatı sonrasında gelişen instabilite nedeni ile uygulanmıştır. Tüm hastalara ameliyat öncesi ve sonrası MRG ve röntgen çekilmiştir. MRG'de komşu segment disk herniasyonu, röntgende kayma ve rotasyonel açılma değerlendirilmiştir.

Bulgular: Üç hastada komşu segmentte disk herniasyonu, 6 hastada instabilite saptandı. İnstabilite ve disk herniasyonu gelişimi ile hastaların yaşı, cinsiyeti, patolojinin tipi veya sabitlenen seviye arasında istatistiksel bağlantı bulunamadı.

Sonuç: Sabitleme ve füzyon komşu segmentte anlamlı derecede instabiliteye neden olmamaktadır. Daha fazla hasta ile ve farklı instabilite ölçütleri ile yapılacak randomize çalışmalar komşu segment instabilitesini daha iyi anlamayı sağlayacaktır.

Anahtar Sözcükler : Dejeneratif lomber omurga hastalığı; enstrümantasyon füzyon; kısa segment füzyon; komşu segment instabilitesi

Kanıt Düzeyi: Retrospektif klinik çalışma, Düzey III

* Specialist of Neurosurgery, Neurosurgery Clinic, Ministry of Health, Diskapi Yıldırım Beyazıt Research and Training Hospital, Ankara;

** Assoc. Prof. of Neurosurgery, Neurosurgery Clinic, Private Kuru Hospital, Ankara;

*** PhD, Assoc. Prof. of Neurosurgery, Neurosurgery Clinic, Private Lokman Hekim Hospital, Ankara;

**** Specialist of Neurosurgery, Neurosurgery Clinic, Ministry of Health, Sevkett Yılmaz Research and Training Hospital, Yıldırım, Bursa.

INTRODUCTION

Stabilization and fusion of the lumbar spine for instability are becoming increasingly common in neurosurgical practice (13). Surgery aims to decompress the spinal cord, stabilize the spine, and correct deformities. Ultimately it prevents chronic spinal deformity and instability. A variety of instruments, bones, and bone-like materials are used for stabilization. The true incidence and clinical impacts of instability at the adjacent upper segment after stabilization of the lumbar spine are unclear (1,3,8-9). This study investigates the development of disc herniation or instability at the adjacent upper segment of the stabilized lumbar spine in 29 patients over an $18,21 \pm 6,42$ month follow-up period (a range of 6-30 months).

MATERIALS AND METHODS

Twenty-nine patients were operated on at the Neurosurgery Clinic, Ministry of Health, Dışkapı Yıldırım Beyazıt Research and Training Hospital, Ankara, Turkey between January 2006 and December due to differing pathologies, these operations used short segment 98 posterior instrumentation and fusion. Patients were evaluated for long-term development of adjacent upper segment disc herniation and instability. Twenty-one patients were male, 8 patients were female, and their mean age at the time of surgery was 51,3 (33 – 69). Stabilization and fusion were performed on L3 - 4 in one patient, on L4-5 in 16 patients, and on L5-S1 in 12 patients. Stabilization and fusion were performed for one-level listhesis or the presumed instability after lumbar stenosis operations, or unstable (radiologic or peroperative) lumbar disc herniation.

In all patients, preoperative lumbar magnetic resonance imaging (MRI), anteroposterior, lateral, hyperflexion, and hyperextension X-ray images were obtained. Anteroposterior and lateral X-ray images were obtained at the first day after surgery. Anteroposterior, lateral, hyperflexion, and hyperextension X-ray images were obtained again six months following surgery. In postoperative lateral X-rays, sagittal slippage and rotational angling of the adjacent upper segments were measured. Scorings were made according to White and Panjabi. In lumbar MRIs, disc herniation and indication of surgery for this disc herniation were evaluated for the adjacent upper disc level. After the lumbar spinal operation, the first follow-up was conducted 1.5 months after the operation. The second follow-up visit was conducted six months after surgery. Patients then returned for follow ups at six-month intervals. Patients included in this study returned for at least

one follow-up, with a maximum of six follow-ups. The mean follow-up period was $18,21 \pm 6,42$ months, with a range of 6-30 months. During each follow-up visit, lumbar MRI, anteroposterior, lateral, hyperflexion, and hyperextension X-ray images were obtained.

In the adjacent upper segment, development of lumbar disc herniation and surgical indication for this disc herniation were evaluated based on the following grading system: Grade 1 (no disc herniation), Grade 2 (non-surgical disc herniation), and Grade 3 (surgical disc herniation). Instability was evaluated according to White and Panjabi's criteria: Grade 1 (no instability; White-Panjabi score 0-2), Grade 2 (limited instability; White-Panjabi score 3-4), Grade 3 (instability; White-Panjabi score greater than or equal to 5).

Statistical analysis:

All statistical analyses were performed using Statistical Package for Social Sciences (SPSS) for Windows (Version 11,5, IBM, Chicago, IL, USA). Continuous variables were shown as mean $\pm$ Standard deviation (minimum-maximum), and categorical variables were shown as case numbers and (%). The effect of one variable on disc herniation and instability was evaluated using Student's t-test. Categorical variables were evaluated using Pearson Chi-square test and Fisher's Chi square exact test. P-values $<0,05$ were considered to be statistically significant.

RESULTS

Patients' demographic data (e.g. age and gender), pathologies, and surgeries are shown in Table-1. Out of 29 patients, 10 patients had listhesis, 5 patients had listhesis plus lumbar disc herniation, and 10 patients had lumbar stenosis plus instability. In postoperative MRIs of the 29 patients, adjacent upper level instability grades and disc herniation grades are shown in Figure 2. One patient had the L2-L3 level as the adjacent upper segment, 17 patients had L3-L4 levels as the adjacent upper segment, and 11 patients had L4-L5 levels as the adjacent upper segment. Twenty-six patients had Grade 1 disc herniation (no disc herniation), 2 patients had Grade 2 disc herniation, and 1 patient had Grade 3 disc herniation. Three patients had two or more grades of disc herniation; two of these patients had stabilization and fusion at L3-4 and one patient had stabilization and fusion at L5-S1. Twenty-three patients had no instability at the adjacent upper segment, while 6 patients had instability at the adjacent upper segment (Table. 1-4).

Table-1. Table showing patients' age, gender, pathology, and level of posterior stabilization.

No	Age	Gender	Diagnose	Level of Posterior stabilization
1	44	F	LISTHESIS	L5-S1
2	43	F	LISTHESIS	L4-5
3	61	F	SPONDILOSIS AND INSTABILITY	L5-S1
4	69	F	SPONDILOSIS AND INSTABILITY	L4-5
5	38	F	LISTHESIS	L5-S1
6	55	F	LISTHESIS+HERNIATED DISC	L4-5
7	41	M	LISTHESIS	L5-S1
8	43	F	LISTHESIS	L5-S1
9	46	M	SPONDILOSIS AND INSTABILITY	L5-S1
10	54	F	SPONDILOSIS AND INSTABILITY	L4-5
11	61	F	SPONDILOSIS AND INSTABILITY	L4-5
12	51	M	LISTHESIS AND HERNIATED DISC	L4-5
13	33	F	LISTHESIS	L4-5
14	69	M	SPONDILOSIS AND INSTABILITY	L4-5
15	59	M	LISTHESIS	L5-S1
16	43	F	LISTHESIS	L5-S1
17	47	F	LISTHESIS AND HERNIATED DISC	L4-5
18	62	F	SPONDILOSIS AND INSTABILITY	L5-S1
19	69	F	SPONDILOSIS AND INSTABILITY	L5-S1
20	35	F	LISTHESIS	L4-5
21	58	F	LISTHESIS AND HERNIATED DISC	L5-S1
22	54	F	LISTHESIS	L4-5
23	54	M	LISTHESIS	L4-5
24	64	F	SPONDILOSIS AND INSTABILITY	L4-5
25	43	F	LISTHESIS	L3-4
26	46	M	LISTHESIS	L4-5
27	44	F	LISTHESIS	L4-5
28	48	M	LISTHESIS AND HERNIATED DISC	L4-5
29	53	F	SPONDILOSIS AND INSTABILITY	L4-5

Table-2. Table showing patients' level of posterior stabilization, adjacent upper segment disc herniation grade and instability grade. In the evaluation of the adjacent upper segment;

No	Level of stabilization	Herniated disc in the adjacent segment*	White &Panjabi Score	Instability in the adjacent segment**
1	L5-S1	1	0	1
2	L4-L5	1	0	1
3	L5-S1	1	3	2
4	L4-L5	1	0	1
5	L5-S1	1	0	1
6	L4-L5	3	3	2
7	L5-S1	2	0	1
8	L5-S1	1	0	1
9	L5-S1	1	3	2
10	L4-L5	1	0	1
11	L4-L5	1	0	1
12	L4-L5	1	0	1
13	L4-L5	1	0	1
14	L4-L5	1	0	1
15	L5-S1	1	0	1
16	L5-S1	1	0	1
17	L4-L5	2	0	1
18	L5-S1	1	3	2
19	L5-S1	1	0	1
20	L4-L5	1	0	1
21	L5-S1	1	0	1
22	L4-L5	1	0	1
23	L4-L5	1	3	2
24	L4-L5	1	0	1
25	L3-L4	1	0	1
26	L4-L5	1	0	1
27	L4-L5	1	3	2
28	L4-L5	1	0	1
29	L4-L5	1	0	1

*1: no disc herniation, 2: non-surgical disc herniation, 3: surgical disc herniation. **1: no instability (White - Panjabi score 0-2), 2: limited instability (White - Panjabi score 3-4), 3: instability (White - Panjabi score greater than or equal to 5).

Table-3. Table showing the relationship between age, gender, pathology, adjacent upper segment of the patients and development of disc herniation at the adjacent upper segment.

Variables	No disc herniation	Disc herniation	p	Variables
Age (years)	51,7 ± 10,5	47,7 ± 7,0	0,526*	Age (years)
Gender			1,000**	Gender
Male	7 (%87,5)	1 (%12,5)		Male
Female	19 (%90,5)	2 (%9,5)		Female
Pathology			0,069***	Pathology
Listhesis	13 (%92,9)	1 (%7,1)		Listhesis
Listhesis + Lumbar disc herniation	3 (%60,0)	2 (%40,0)		Listhesis + Lumbar disc herniation
Stenosis + Instability	10 (%100)	-		Stenosis + Instability
Adjacent segment			0,872***	Adjacent segment
L2-L3	1 (%100)	-		L2-L3
L3-L4	15 (%88,2)	2 (%11,8)		L3-L4
L4-L5	10 (%90,9)	1 (%9,1)		L4-L5

*Student's t-test, **Fisher's Chi-square exact test, ***Pearson's Chi-square test.

Table-4. Table showing the relationship between age, gender, pathology, the adjacent segment of the patients and development of instability at the adjacent segment.

Variables	No instability	Instability	p
Age (years)	50,3 ± 10,3	55,0 ± 9,5	0,323*
Gender			1,000**
Male	6 (%75,0)	2 (%25,0)	
Female	17 (%81,0)	4 (%19,0)	
Pathology			0,649***
Listhesis	12 (%85,7)	2 (%14,3)	
Listhesis + Lumbar disc herniation	4 (%80,0)	1 (%20,0)	
Stenosis + Instability	7 (%70,0)	3 (%30,0)	
Adjacent segment			0,745***
L2-L3	1 (%100)	-	
L3-L4	13 (%76,5)	4 (%23,5)	
L4-L5	9 (%81,8)	2 (%18,2)	

*Student's t-test, **Fisher's Chi-square exact test, ***Pearson's Chi-square test.

Herniation at the adjacent upper segment did not correlate with age ($p=0,526$), gender ($p=1,000$), pathology type ($p=0,069$), and level of stabilization ($p=0,872$). Development of instability at the adjacent upper segment did not correlate with age ($p=0,323$), gender ($p=1,000$), pathology type ($p=0,649$), and level of stabilization ($p=0,745$).

DISCUSSION

Instability, which can cause pain and neurologic deficits, is a major problem for Spine surgeons. Surgery for instability is becoming more common as surgeons develop new techniques. However, stabilization and fusion of unstable segments prevent normal physiologic movement, change the biomechanics of the vertebral column, and provoke adjacent segment degeneration (2,6,7,12). Researchers are currently 155 developing ways to reduce these biomechanical changes and preserve normal physiologic movement. New surgical techniques, treatment models, and surgical instruments are being developed. Stabilization and

fusion surgeries are essential for the treatment of listhesis, acute traumatic vertebra fractures, and degenerative disc diseases. However, sometimes pain is not caused by a loss of stability in the vertebral column. Instead pain may be caused by the loss of distribution of load. In these cases dynamic stabilization is necessary (10). Lumbar listhesis is common with spinal disc diseases and spinal stenosis in elderly patients. In these cases, neurologic decompression should be used. In patients with symptomatic stenosis, facetectomy

is performed during surgery. This procedure causes instability and related complaints. Various stabilization techniques are used to treat lumbar fractures, degenerated discs, and tumoral diseases. Anterior, posterior, and combined techniques are used for stabilization (15). Each technique has advantages and disadvantages. The technique used depends on the patient's clinic and pathology, as well as the surgeon's experience.

Suk et al. (14) operated on 76 patients with degenerative spondylolisthesis. The researchers performed decompression, transpedicular stabilization, and posterolateral fusion on 36 patients. Suk et al. also performed interbody fusion on 40 patients. In a two-year follow-up, the first group showed 92% fusion and the second group showed 100% fusion. In first group there was 28% listhesis, while in second group there was 45% listhesis. The researchers concluded that a large fusion area was needed in order to achieve fusion over a long-term follow-up period.

One of the aims of posterior stabilization and fusion surgery is to obtain normal lumbar lordosis. Putting cages between vertebral bodies helps in obtaining normal lumbar lordosis. Godde et al. (4) compared rectangular and wedge-shaped cages. They 178 found that lumbar lordosis increased in patients with wedge-shaped cages and decreased in patients with rectangular cages. A decrease in lumbar lordosis has a negative effect on biomechanics and increases mobility in the adjacent upper segment. Studies have shown that fusion rates were higher in patients who underwent stabilization and fusion. However, adjacent segment diseases increased after rigid fixation (11). Hillibrand and Robbins (5) defined adjacent segment diseases as new symptoms, i.e. radiographic changes that developed after spinal fusion surgery. In these patients symptoms improved after surgery, but new symptoms eventually developed because of adjacent segment diseases.

In this study we evaluated the stabilized and fused adjacent upper segments of 29 patients. Three patients developed disc herniation in the adjacent upper segment, but only one of them needed surgery. None of the patients developed instability requiring surgery at the adjacent upper segment. However, 6 patients developed instability not requiring surgery. Patients who had surgery because of disc herniation were also stabilized.

A pain-free spine capable of normal physiologic movement is important for any patient's quality of life. Treatment of spine pathologies should consider anatomic and physiologic rules. Surgical techniques should not harm normal structures. Stabilized vertebrae should be as short as possible, and the stabilization material should be suitable for the anatomy; it should be strong and allow physiologic movement. After surgery, patients should be monitored for contractures and loss of strength in the muscles surrounding spine. Patients should be given physical therapy to strengthen these muscles and to relieve contractures.

There is a burgeoning interest among surgeons in increased movement and loading at the adjacent segment after stabilization and the development of adjacent segment disease. Adjacent segment disease is defined as radiologic degenerative changes and new symptoms known to occur following reconstructive spine surgery, most commonly immediately adjacent to the functional spinal unit.

In this study, the development of adjacent upper segment instability and disc herniation was evaluated in patients who underwent stabilization and fusion operations for different pathologies. The study included a sufficient follow-up period. The study found that adjacent upper segment instability was not very common. However, some patients did require surgery. Our data showed that rigid stabilization and fusion were sufficient for the patients. These patients did not display too much instability and disc herniation at the adjacent upper segment. Nevertheless, it is important to remember that every patient is different. Additional randomized studies with more patients and different instability criteria are needed to understand adjacent upper segment instability. Minimally invasive approaches and surgeries respecting normal physiologic angles should help to obtain optimum results.

REFERENCES

- 1- Aiki H, Ohwada O, Kobayashi H, Hayakawa M, Kawaguchi S, Takebayashi T, Yamashita T. Adjacent segment stenosis after lumbar fusion requiring second operation. *J Orthop Sci* 2005; 10: 490-495.
- 2- Cakir B, Carazzo C, Schmidt R, Mattes T, Reichel H, Käfer W. Adjacent segment mobility after rigid and semi-rigid instrumentation of the lumbar spine. *Spine* 2009; 34: 1287-1291.
- 3- Cheh G, Bridwell KH, Lenke LG, Buchowski JM, Daubs MD, Kim Y, Baldus C. Adjacent segment disease following lumbar/thoracolumbar fusion with pedicle screw instrumentation: a minimum 5-year follow up. *Spine* 2007; 32: 2253-2257.
- 4- Godde S, Fritsch E, Dienst M, Kohn D. Influence of cage geometry on sagittal alignment in instrumented posterior lumbar interbody fusion. *Spine* 2003; 28: 1693-1699.
- 5- Hilibrand AS, Robbins M. Adjacent segment degeneration and adjacent segment disease: the consequences of spinal fusion? *Spine J* 2004; 4: 190-194.
- 6- Hirabayashi K, Maruyama T, Wakano K, Ikeda K, Ishii Y. Postoperative lumbar canal stenosis due to anterior spinal fusion. *Keio J Med* 1981; 30: 133-139.
- 7- Lee CK, Langrana NA. Lumbosacral spinal fusion. A biomechanical study. *Spine* 1984; 9: 574-581.
- 8- Levin DA, Hale JJ, Bendo JA. Adjacent segment degeneration following spinal fusion for degenerative disc disease. *Bull NYU Hosp Jt Dis* 2007; 65: 29-36.
- 9- McCormick PC. The adjacent segment. *J Neurosurg Spine* 2007; 6: 1-4.
- 10- Mulholland RC, Sengupta DK. Rationale, principles and experimental evaluation of the concept of soft stabilization. *Eur Spine J* 2002; 11: 198-205.
- 11- Resnick DK, Weller SJ, Benzel EC. Biomechanics of the thoracolumbar spine. *Neurosurg Clin N Am* 1997; 8: 455-469.
- 12- Sears WR, Sergides IG, Kazemi N, Smith M, White GJ, Osburg B. Incidence and prevalence of surgery at segments adjacent to a previous posterior lumbar arthrodesis. *Spine J* 2011; 11: 11-20.
- 13- Sonntag VK, Marciano FF. Is fusion indicated for lumbar spinal disorders. *Spine* 1995; 20: 138-142.
- 14- Suk SI, Lee CK, Kim WJ, Lee JH, Cho KJ, Kim HG. Adding posterior lumbar interbody fusion to pedicle screw fixation and posterolateral fusion after decompression in spondylolytic spodylolisthesis. *Spine* 1997; 22: 210-219.
- 15- Zhang JW, Xiao BP, Xu RM, Zhao LJ, Ma WH, Ruan YP. Analysis of safety and effect of reconstructing anterior and middle columns by single posterior approach in treating lumbar burst fractures. *Chin J Traumatol* 2009; 12: 107-112.

Adress: Uygur ER, Söğütözü Caddesi, 4. Sokak, No: 22/7, 06510, Cankaya, Ankara, Turkey
Tel: 0505 589 23 55
Fax: 0312 284 51 02
E-mail: uygurer@gmail.com
Geliş tarihi: 11 Aralık 2012
Kabul tarihi: 16 Ocak 2013