

Comparison Between Carpal Tunnel Syndrome and Individuals With Symptoms Of Obsessive-Compulsive Disorder

Karpal Tünel Sendromu ile Obsesif Kompulsif Hastalık Semptomları Arasındaki İlişki

Meliha GÜNDAĞ PAPAKER*0000-0003-1271-9023

Anas ABDALLAH* 0000-0003-3600-089x

M.Hakan SEYİTHANOĞLU* 0000-0002-6072-0347

Engin CAN* 0000-0002-1343-0443

Yıldızhan ŞENGÜL** 0000-0002-5087-9944

Aygül TANTİK PAK** 0000-0002-7414-3800

*Bezmialem Vakıf Üniversitesi Tıp Fakültesi
Hastanesi, Nöroşirurji Anabilim Dalı, İstanbul,
Türkiye

**Gaziosmanpaşa Taksim Eğitim ve Araştırma
Hastanesi, Nöroloji Kliniği, İstanbul, Türkiye

Yazışma Adresi: Meliha GÜNDAĞ PAPAKER

Bezmialem Vakıf Üniversitesi Tıp Fakültesi Hastanesi

Nöroşirurji Anabilim Dalı, İstanbul, Türkiye

E-Mail: melihagundag34@gmail.com

Öz

Amaç: Karpal Tünel Sendromu (KTS) en sık görülen periferik sinir tuzak nöropatisidir ve genel popülasyonun %3'ünde görülür. Bu çalışmanın amacı, KTS'li hastalarda Moudsley Obsesif Kompulsif Envanteri (MOKE) ile temizlik kompulsiyonu semptomları ile KTS arasındaki ilişkiyi incelemektir.

Metod: Bezmialem Vakıf Üniversitesi Hastanesi Nöroşirurji Polikliniğine başvuran 60 KTS'li hasta ile 60 sağlıklı kontrol grubu değerlendirildi. Nörolojik muayene, elektrofizyolojik bulgular, Leeds assessment of neuropathic symptoms and signs (LANSS) ile nöropatik ağrı varlığı, MOKE ile Obsesif Kompulsif Hastalık belirtileri, Beck Anksiyete Envanteri (BAE) ile anksiyete düzeyleri değerlendirildi.

Bulgular: Anksiyete düzeyleri ile KTS arasında anlamlı bir ilişki gözlenmedi ($p > 0.005$). MOKE ruminasyon, kontrol ve şüphe alt ölçekleri ile KTS arasında anlamlı bir ilişki gözlenmedi ($p > 0.005$). Hasta grubu kontrol grubuna göre daha yüksek MOKE temizlik alt ölçeği puanına sahipti ve fark istatistiksel olarak anlamlıydı ($p = 0.008$). Hasta grubu kontrol grubuna göre daha yüksek MOKE yavaşlık alt ölçeği puanına sahipti ve fark istatistiksel olarak anlamlıydı ($p = 0.004$).

Sonuç: Temizlik takıntısı bulguları olan bireylerde artan el bilek hareketleri, KTS oluşumu riski ile ilişkilidir. Çalışmamız çok fazla seviyede olan el bilek fleksiyon/ekstansiyon hareketlerinde azalmanın KTS vakalarını azaltabileceğini göstermiştir.

Anahtar Kelimeler: Karpal Tünel Sendromu, LANSS, Obsesif Kompulsif Hastalık

Abstract

Objective: Carpal tunnel syndrome (CTS) is the most common peripheral nerve entrapment neuropathy and is seen in 3 % of the general population. The aim to this study was examine the relationship between the symptoms of cleaning compulsion with using Moudsley obsessive compulsive inventory (MOCI) and patients with CTS.

Methods: Sixty patients with CTS and 60 healthy controls were evaluated in Bezmialem Vakıf University Faculty of Medicine Neurosurgery Clinic. Neurological examination, Electrophysiological findings, the presence of neuropathic pain evaluated by the Leeds assessment of neuropathic symptoms and signs (LANSS) , the symptoms of obsessive- compulsive disorder using with MOCI and anxiety levels assessed by the Beck Anxiety Inventory (BAI) were evaluated.

Results: No statistically significant relationship was observed between anxiety levels and CTS ($p > 0.005$). No statistically significant relationship was found MOCI rumination, control and doubting subscale scores ($p > 0.005$).

Geliş Tarihi: 16/02/2021

Kabul Tarihi: 20/10/2021

Patient group had higher MOCI cleaning subscale scores than the control group and the difference was statistically significant ($p = 0.008$). Patient group had higher MOCI slowness subscale scores than the control group and the difference was statistically significant ($p = 0.004$).

Conclusion: Increased wrist movement in people that have the symptoms of compulsive cleaning is associated with the risk of CTS. Our study show that a reduction of high level of wrist flexion-extension movements may decrease the new CTS cases.

Keywords: Carpal tunnel syndrome, LANSS, Obsessive compulsive disorder

Introduction

Carpal tunnel syndrome (CTS) is the most common peripheral nerve entrapment and is seen in 3 % of the general population (1). The carpal tunnel (CT) is formed by the wrist bones, the transvers ligament (TL), flexor retinaculum, and the median nerve (MN) passes into the CT (2). Due to its specific location in CT, MN is highly susceptible to compression, and in repeated wrist flexion/extension movement, the pressure in CT increases and compresses MN and symptoms begin (2). Typical symptoms include numbness, paraesthesia and pain in the areas innervated by MN, weakness and atrophy of the thenar muscles which affects the quality of life and work efficiency in affected patient (1,6). Electromyography (EMG) is 49-84 % sensitive and 95-99 % specific for diagnosis (3). The Leeds Assessment of Neuropathic Symptoms and Signs (LANSS) scale is a clinical-based test used to identify patients whose pain is dominated by neuropathic mechanisms (4).

People who have the symptoms of Obsessive compulsive disorder (OCD) feels need to perform certain routines repeatedly called compulsion and thoughts repeatedly called obsessions and they don't control either the activities and thoughts in a short period of time (5). Especially in individuals with symptoms of cleaning activities, due to excessive flexion/extension movements MN could be trapped in CT, so between these persons and CTS was evaluated. Psychiatric disorders can also be observed in people have the symptoms of OCD, anxiety levels were also evaluated in our study (6).

Methods

Our study was carried out Bezmialem Vakif University Faculty of Medicine Neurosurgery Clinic, consist of 60 CTS patients and 60 control groups, between January 2017 and March 2019. Patients with older than 18 and younger than 65 years old were included in this study. The exclusion criteria were older than 65 years old, cervical disk herniation and cervical canal stenosis. EMG findings of upper extremity has two groups minimal-mild and moderate-severe CTS.

Neuropathic pain was assessed with LANSS that consist of two parts pain questionnaire and sensation. Pain questionnaire consist of 5 questions and is worth 16 points. And a total of 8 points including 5 points allodynia and 3 points pin-prick threshold sections. In this way, the LANSS test is totally 24 points. Twelve and below not have neuropathic pain between 13 and 24 have neuropathic pain (8).

Beck anxiety inventory (BAI) is required to measure standard anxiety levels. The BAI consist of 21 questions each has between 0-3. Between 0-9 shows minimal, 8-15 mild, 16-25 medium and 26-63 serious anxiety (10).

Mouldsley obsessive compulsive inventory (MOCI) is a self report scale performed by Hodgson and Rachman and used to measure the type and severity of OCD symptoms (10,22). The MOCI scale is consist of 30 items and 4 sub-scales that are called checking, cleaning, slowness and doubting. Adaptation of the scale into Turkish, validity and reliability of the Turkish form were done by Erol and Savasir (9). Turkish form of MOCI contains 37 items (7 items from Minnesota Multiple Personality Inventory [MMPI]) and 5 sub-scales (5th sub-scale is called rumination). The measurement is made according to the "right" or "wrong" responses given by the participants. As the MOCI scale scores increase, it is assumed that the severity of OCD symptoms increases (7).

This study was approved by Bezmialem Vakif University noninterventional research Ethics Committee (18.07.2018. 14/14).

Statistical Analysis

Statistical analysis of the data of our study which was planned as a cross-sectional study, was performed using SPSS 24.0 package program. All numerical variables were expressed as mean $\pm$ standard deviation, and categorical variables were represented by frequency and probability tables. Kolmogorov Smirnov and Shapiro-Wilk tests were performed to test the compatibility of the sample with normal distribution in terms of age and educational level. It was shown that it had normal distribution. Due to the compatibility with the normal distribution, independent samples t-test (Student t-test) was used for comparison of numerical data and the chi-square test was used for comparison of categorical variables. Pearson correlation test was applied to investigate the correlation between numerical data. A p value < 0.05 was accepted as statistically significant in all statistical analyses.

Results

A total of 60 CTS patients, 54 (90%) women and 6 (10%) men were included in our study. The mean age of patients was 46, (26 ± 8.76 years). Eight patients have hypertension, 6 have diabetes mellitus, 2 have hypothyroidism, 1 has OCD, 1 has pregnant and 1 has cow milking work for a long years. There was no comorbidity in 41 of the patients. According to the EMG findings, patients were analyzed in 2 groups minimal-mild and moderate-severe CTS.

Twenty-eight of the patients were minimal to mild and 32 were moderate to severe CTS. Neuropathic pain was present in 33 (55%) of the patients. Four of the patients had surgery. According to the BAI, 15 (25%) patients have minimal, 17 (28%) mild, 16 (27%) moderate and 12 (20%) severe anxiety syptoms. No significant relationship was observed between anxiety levels and CTS ($p > 0.005$). No significant relationship was observed between MOCI (checkingsub-scale)and CTS ($p > 0.005$). No significant relationship was observed between MOCI (doubtingsub-scale) and CTS ($p > 0.005$). A significant relationship was observed between MOCI (cleaningsub-scale) and CTS ($p = 0.008$). A significant relationship was observed between MOCI (slownesssub-scale) and CTS ($p = 0.004$) Table 1.

Table1: Anxiety and Moudsley obsessive compulsive inventory scales of investigated 60 patients with Carpal Tunnel Syndrome.

	group	n	mean	Standard deviatio n	F	t	p
Age	CTS	60	46.3	8.8	18.5	1.8	0.068
	control	60	42.6	13.0		1.8	0.069
Anxiety	CTS	59	1.3	1.1	1.3	-.2	0.871
	control	60	1.4	1.8		-.2	0.870
MOCI(controlsub-scale)	CTS	50	2.9	2.0	0.3	.1	0.922
	control	58	2.9	2.2		.1	0.922
MOCI(cleaningsub-scale)	CTS	50	4.8	3.0	25.8	2.7	0.008
	control	58	3.6	1.5		2.6	0.011
MOCI(slownesssub-scale)	CTS	50	2.5	1.5	3.9	3.0	0.004
	control	58	1.7	1.2		2.9	0.004
MOCI(doubtingsub-scale)	CTS	50	3.2	1.7	2.7	.2	0.874
	control	58	3.2	1.3		.2	0.877
MOCI(ruminationsub-scale)	CTS	50	2.2	5.2	1.4	-.5	0.599
	control	58	3.7	20.1		-.6	0.576

Discussion

Most patients of CTS are manifested by pain and numbness. Understanding the mechanism of pain is necessary to create an appropriate treatment plan. The LANSS pain scale is a validated tool for distinguishing patient with neuropathic pain from with nociceptif pain. Gursoy et al. showed that neuropathic pain mechanisms in CTS does not have a significant relationship with CTS severity in EMG (8). In another study, Truini et al. reported that there was no difference between the the presence of neuropathic pain and CTS severity in 70 CTS patients (14). In our study 28 patients had minimal to mild and 32 patients had moderate to severe CTS. We found neuropathic pain in 33 (55%) of our patients.

Lund et al. Measured wrist angular velocity, mean power frequency and range of motion on 33 different professions. They reported that statistically significant relationship between wrist movements level with risk of CTS (7). In another study, Violente et al found an increased incidence of CTS for workers exposure to fast and forceful manual work (11). Mattioli et al. reported that prolonged flexion-extension movement of the wrist, and especially the use of vibrating hand tools, increase the risk of CTS (12). In our study , we found that statistically significant relationship between individuals with symptoms of cleaning compulsion and the risk of CTS ($p=0.008$).

CTS usually occurs with repetitive movements of wrist and resulting in synovial sheath hypertrophy, tenosynovitis, intracarpal swelling and MN compression (1,15,16). Rheumatoid arthritis, scleroderma, systemic lupus erythematosus, polymyalgia rheumatica, collagen tissue diseases, diabetes mellitus, pregnancy, estrogen and oral contraceptive use, acromegaly, myxedema, hyperparathyroidism, amyloidosis, tumors such as lipoma and ganglioma, chronic trauma are diseases and conditions associated with CTS. However usually the cause of CTS is unknown called idiopathic CTS (17).

The treatment is carried out according to the cause of the CTS. For example, it is thought that during pregnancy CTS is caused by fluid retention, and postpartum resolves spontaneously (18). Conservatively, a splint is recommended to holding the wrist at 15 degress of dorsiflexion to reduce the night awakening of the patients. In addition, non-steroidal anti-inflammatory drugs ,physical therapy methods such as ultrasound and electrical nerve stimulation , can also be used. If the CTS symptoms do not improve with treatment of systemic disease, if conservative treatment methods are not effective and the symptom are progressing, if there is neurological motor loss, EMG nerve conduction abnormalities, surgical decompression should be considered.In surgery, it is aimed to remove the median nerve compression by cutting the transverse ligament completely. Surgical intervention may fail due to insufficient incision of the transverse ligament, cutting of the MN palmar motor branch, and the development of abnormal granulation. In this case, symptoms may continue in the same way, recur after a certain improvement, and new symptoms may develop (19).

As surgical treatment methods, open surgery, microsurgery or endoscopic surgery methods can be used. In generally local anesthesia is used in open surgery. In open surgery, which is a minimally invasive technique, standard incision and mini incision can be applied. The longitudinal long incision is used in the standard incision, the advantage is that all carpal tunnel elements are visible, and the disadvantage is that the scar tissue is large and can lead to loss of hand function. Mini incision is preferred due to minimal scar tissue but since all of the carpal tunnel elements cannot be seen, it is necessary to check whether the transverse ligament is completely cut. The advantage of standard open surgery over all other surgical methods is to dissect and protect the MN motor branch. There are those who claim that microsurgery is the most important advantage of fully viewing and incising the distal part of the transverse ligament, as well as those who claim the same advantage in open surgery.

In a study comparing endoscopic surgery with open surgery, it was reported that infection, hypertrophic scar tissue and scar sensitivity are more common in open surgery, and when total number of complications is evaluated, endoscopic surgery is more reliable.

Also, in endoscopic surgery, the muscle and skin on the transverse carpal ligament are preserved, providing better grip strength and earlier return to work. However, the most important disadvantages of endoscopic surgery are injury to MN and superficial palmar arcus. As a result, regardless of the surgical method, recovery rates in CTS are reported to above 90 % (20,21).

Conclusion

Our study is associated with an increased risk of CTS due to recurring wrist movements in individuals with symptom of cleaning compulsion. The management of reducing the repetitive movement in the wrist of the individuals with symptom of cleaning compulsion shows that the development of new CTS cases can be prevented. Our study is retrospective, single-center and limited, since it contains few cases. However, large-scale studies are needed.

References

1. Trull-Ahuir C, Sala D, Chismol-Abad J, Vila-Caballer M, Lion Jf. Efficacy of platelet-rich plasma as an adjuvant to surgical carpal ligament release: a prospective, randomized controlled clinical trial. *SciRep*. 2020 Feb 7;10(1):2085. doi: 10.1038/s41598-020-59113-0
2. Rayegani SM, Moradi-Joo M, Raeissadat SA, Bahrami MH, Seyed-Nezhad M, Heidari S. Effectiveness of Low-Level Laser Therapy compared to Ultrasound in Patients With Carpal Tunnel Syndrome: A Systematic Review and Meta-analysis. *J Lasers Med Sci*. 2019 Fall;10(Suppl 1):S82-S89. doi: 10.15171/jlms.2019.S15. Epub 2019 Dec 1.
3. Simsir Atalay N, Sarsan A, Akkaya N, Yıldız N, Topuz O. The Impact of Disease Severity in Carpal Tunnel Syndrome on Grip Strength, Pinch Strength, Fine Motor Skill and Depression J. *Phys. Ther. Sci*. 23: 115–118, 2011
4. Bennett M. The LANSS Pain Scale: the Leeds Assessment of Neuropathic Symptoms And Signs. *Pain*. 2001;92:147–157
5. Hu Y, Liao R, Chen W, Kong X, Liu J, Liu D, Maguire P, Zhou S, Wang D. Investigating behavior inhibition in obsessive-compulsive disorder: Evidence from eye movements. *Scand J Psychol*. 2020 Feb 6. doi: 10.1111/sjop.12620. [Epub ahead of print] PMID: 32027033
6. Padua L, Coraci D, Erra C, Pazzaglia C, Paolasso I, Loreti C, Caliendo P, Hobson-Webb LD. Carpal tunnel syndrome: clinical features, diagnosis, and management: *Lancet Neurol* 2016; 15: 1273–84
7. Lund CB, Mikkelsen S, Thygesen LC, Hansson G, Thomsen JF. Movements of the Wrist and the Risk of Carpal Tunnel Syndrome: A Nation wide Cohort Study Using Objective Exposure Measurements: *Occup Environ Med* 2019 Aug;76(8):519-526. doi: 10.1136/oemed-2018-105619. Epub 2019 Jun 12.
8. Gürsoy AE, Kolkusa M, Babacan Yıldız G, Kocaman G, Çelebi A, Koçer A: Relationship between electrodiagnostic severity and neuropathic pain assessed by the LANSS pain scale in carpal tunnel syndrome. *Neuropsychiatr Dis Treat*. 2013; 9: 65–71.
9. Erol N, Savaşır İ. Maudsley Obsessif-Kompulsif Soru Listesinin Türkiye Uyarlaması. In: XXIV Ulusal Psikiyatri ve Nörolojik Bilimler Kongresi Bildiri Kitabı. Ankara: Gata Basımevi: 1988.p.107-14.
10. Şengül Y, Şengül HS, Gökçal E, Üstün İ, Öztürk A, Yılmaz O, Babacan Yıldız G. Esansiyel Tremor ve Aleksitimi *Türk J Neurol* 2018;24:248-251.
11. Violante FS, Farioli A, Graziosi F, et al. Carpal tunnel syndrome and manual work: the OCTOPUS cohort, results of a ten-year longitudinal study. *Scand J Work Environ Health* 2016; 42: 280-90.
12. Mattioli S, Baldasseroni A, Curti S, Cooke RM, Mandes A, Zanardi F, et al. Incidence rates of surgically treated idiopathic carpal tunnel syndrome in blue- and white-collar worker and housewives in Tuscany, Italy. *Occup Environ Med*. 2009;66(5):299–304.
13. Abbas MA, Afifi AA, Zhang ZW, et al. Meta-analysis of published studies of work related carpal tunnel syndrome. *Int J Occup Environ Health* 1998; 4: 160-7.
14. Truini A, Padua L, Biasiotto A, et al. Differential involvement of A-delta and A-beta fibres in neuropathic pain related to carpal tunnel syndrome. *Pain*. 2009;145(1–2):105–109.
15. Saint-Lary, O., Rébois, A., Mediouni, Z. & Descatha, A. Carpal Tunnel Syndrome: Primary Care and Occupational Factors. *Front. Med*. 2, (2015).
16. Aboonq, M. S. Pathophysiology of carpal tunnel syndrome. *Neurosciences (Riyadh)*. 20, 4–9 (2015).
17. Harrison MJG: Lack of evidence of generalized sensory neuropathy in patients with carpal tunnel syndrome. *J. Neurol Neurosurg Psychiatry* 1978; 41: 957-959
18. Aminoff MJ. Pregnancy and Disorders of the Nervous System. In: Aminoff MJ: *Neurology and General Medicine*. 2nd ed USA: Churchill Livingstone Inc : 567-585, 1995
19. Ulvi H, Yiğiter R, Aygül R, ÖV Asuman: Karpal tünel sendromunda tanı ve tedavi yaklaşımları. *Van Tıp Dergisi*: 11 (4):155-160, 2004
20. Vasiliadis HS, Georkoulas P, Shrier I, Salanti G, Scholten RJPM: Endoscopic release for carpal tunnel syndrome. *Cochrane Database Syst Rev*. 2014 Jan 31; (1): CD008265
21. Açıkgöz B, Sümer M: Karpal Tünel Sendromu. *Türk Nöroşirurji Dergisi* 10: 79-84, 2000
22. Hodgson RJ, Rachman S. Obsessional-compulsive complaints. *Behav Res Ther* 1977;15:389-95.