

PREDICTORS OF IMPROVEMENT IN LEFT VENTRICULAR FUNCTION AFTER STENT IMPLANTATION OF CHRONIC CORONARY OCCLUSION

¹MD. Ferit Akgül, ²MD. Talantbek Batyraliev, ²MD. Zarema Karben, ²MD. Alper Serçelik, ²MD. Aytaç Vural, ²MD. Denis Fettser

¹Mustafa Kemal University, Faculty of Medicine, ANTAKYA, ²Sani Konukoglu Medical Center, Cardiology Department, GAZİANTEP

Kronik koroner oklüzyonlu hastalarda sol ventrikül (SIV) performansı üzerine koroner stent implantasyonunun etkisi iyi bilinmemektedir. Bu çalışmanın amacı, bare metal ve ilaç salan stent implantasyonunun SIV ejeksiyon fraksiyonu (SIVEF) etkisini ve SIVEF düzelmesinde hangi klinik ve anjiyografik faktörlerin etkili olabileceğini araştırmaktır. Çalışmamıza en az bir aydır mevcut major bir koroner arter kronik oklüzyonu için başarılı stent implante edilmiş 304 hasta alındı. Ekokardiyografik SIV incelemeleri stent implantasyonu öncesinde ve 6 ay sonra yapıldı. Kronik koroner oklüzyona stent implantasyonundan sonra hem diyastol sonu volüm indeksinde (85.6 ± 18.9 ml/m² ve 80.1 ± 17.1 ml/m²; $p < 0.001$) ve hem de sistolik volüm indeksinde (40.0 ± 15.8 ml/m² ve 34.1 ± 14.3 ml/m²; $p < 0.0001$) azalma ile birlikte SIVEF de ise önemli artış ($53.2 \pm 11.9\%$ ve

$57.0 \pm 11.1\%$; $p < 0.0001$) tüm gruplarda saptandı. Bare metal stent ile ilaç salan stent grupları arasında SIVEF de artış yönünden anlamlı fark yoktu. Multivariate analiz, bazal SIVEF %50 veya düşük, 2 aydan kısa oklüzyon süresi ve diyabetes mellitusun SIVEF de düzelmenin bağımsız belirleyicisi olduğunu ortaya koydu.

Sonuç olarak, kronik koroner oklüzyona stent implantasyonu, ilk 6 ayda, SIVEF üzerinde olumlu etki yapmıştır. Bu olumlu etki özellikle sol ventrikül fonksiyonu deprese olan ve 2 ay ve daha kısa süreli oklüzyonu bulunan hastalarda elde edilmiştir.

Anahtar kelimeler: Kronik koroner oklüzyon, Stent implantasyonu, Sol ventrikül fonksiyonu

(Türk Girişimsel Kard. Der. 2009;13:123-130)

INTRODUCTION

Left ventricular ejection fraction (LVEF) is a powerful predictor of prognosis in patients with coronary artery disease^{1,2}. Interventions that increase LVEF reduce in-hospital mortality and improve long term clinical outcomes^{1,3}. There have been many reports describing late improvement in left ventricular (LV) function following early restoration of coronary artery patency after acute myocardial infarction (MI), but there are only a few studies assessing changes in LV function after revascularization of chronic total coronary occlusions (CTO)⁴⁻¹³. Percutaneous transluminal coronary angioplasty (PTCA) of CTO can be performed with a

good success rate. However, the long-term improvement of LVEF after PTCA for CTO is hampered by high rates of reocclusion and restenosis^{11,12,14-16}. Technological advances in interventional cardiology and the use of coronary stent implantation in patients with CTO results in a much greater increase in lumen diameter and reduces angiographic restenosis compared with PTCA alone^{17,18}. There are limited datas on recovery of LV function after stent implantation in patients with CTO [7-10]. Moreover, the predictors of recovery of LV function after the stent implantation in this group of patients is not well known. So, the aim of this study was to assess the effect of coronary stent implantation on LVEF and to examine what clinical and angiographic factors may have an effect on recovery of LVEF.

METHODS

Study population: The inclusion of patients took place between 7 April 1998 and 24 February 2006 in Sani

Correspondence Address: Dr. Ferit AKGÜL
Konarlı Mah. Sezer Sitesi No:42
Karaağaç/İskenderun/HATAY
Tel: +90 326 214 06 49
Fax: +90 326 614 38 34
E-mail: feritakgull@yahoo.ca
Geliş Tarihi:17.09. 2009
Kabul Tarihi:08.08.2009

Table 1: Clinical and angiographic characteristics of patients underwent stent implantation

Baseline characteristics (n= 304)	
Age (years)	60.4±10.8
Male patients (%)	87.9
Cigarette smoking (%)	
Current	41.1
Past	42.8
Diabetic mellitus (%)	19.4
Treated hypertension (%)	44.0
Family history of CAD (%)	36.8
History of myocardial infarction (%)	54.9
Previous PTCA (%)	14.5
Previous CABG operation (%)	0
Duration of occlusion (months)	6.19±7.0
CCS class	
1, 2	37.1
3, 4	46.7
Medication at discharge	
Antiplatelet	98.0
ACE inhibitor (%)	36.2
Nitrate (%)	58.5
Beta-blocker (%)	61.8
Calcium antagonist (%)	29.6
Statin (%)	46.0
Collateral circulation (%)	60.5
Target-lesion coronary artery (%)	
Left anterior descending	40.8
Left circumflex	27.3
Right coronary	31.9
Stents per patient (%)	
1	77.3
2	21.7
3	1
Stent type (%)	
Bare Metal Stent	54.3
Drug-Eluting Stent	45.7

CAD: coronary artery disease, PTCA: percutaneous transluminal coronary angioplasty, CABG: coronary artery bypass graft surgery, CCS: Canadian Cardiovascular Society, ACE: angiotensin-converting enzyme.

Konukoğlu Medical Center (Gaziantep-Turkey). During this time, 82615 patients underwent diagnostic coronary angiography. Among these patients, 447 patients who had CTO and angina pectoris or exercise-induced ischaemia related to the occluded artery were included in this study.

A CTO was defined as complete interruption of the vessel with Thrombolysis In Myocardial Infarction flow grade 0 that was >1 months old based on the clinical history¹⁴. Occlusions in patients in whom the clinical history did not allow determination of the age of the occlusion were classified as chronic. Patients

who had a MI <1 month of the procedure and significant left main coronary artery disease, patients who had an estimated CTO duration <1 month were excluded. The study was approved by the local Ethics Committee and written informed consent was obtained from all patients before the procedure.

Angiography: Coronary angiography and stenting were performed according to ACC/AHA guidelines for percutaneous coronary intervention¹⁹. Of 447 patients with CTO, 384 patients underwent successful drug-eluting (DES) or bare metal coronary stent (BMS) implantation. The remaining 63 patients who had unsuccessful stent implantation, 34 patients underwent CABG (coronary artery bypass graft) surgery and 29 patients had medical treatment alone. Among the patients who underwent successful stent implantation, 214 patients underwent BMS implantation (Medtronic S670, Medtronic Vascular, Santa Rosa, California; Multi-Link Tristar, Guidant, Temecula, California; BX Velocity, Cordis Corp and Ephesos, Nemed Ltd, Turkey) and 170 patients underwent DES implantation (Cypher, Cordis, Johnson and Johnson Co, The Netherlands and Taxus, Boston Scientific, Boston, Massachusetts). Premedication consisted of 300 mg/day of aspirin orally and 150 mg of clopidogrel starting >24 hours before percutaneous coronary intervention. A bolus of 10,000 IU of heparin was administered after sheath insertion, with repeat bolus given as needed to maintain an activated clotting time of >250 seconds. After the procedure, aspirin 300 mg/day was continued indefinitely and clopidogrel 75 mg/day was administered for 1-6 months.

Echocardiography: Baseline echocardiography (Acuson CV70, Siemens Medical Solutions USA, Inc) was performed before percutaneous intervention. Estimates of LV end-diastolic volume index (EDVI), end-systolic volume index (ESVI), and ejection fraction (EF) were obtained from the average of three consecutive cardiac cycles taken from apical four chamber views using the modified Simpson's rule.

Follow-up: All patients were asked to return for a 6-months follow-up control, regardless of symptomatic status. Within 6 months after the stent implantation, 5 patients died, 16 patients underwent CABG surgery and 36 patients had repeat percutaneous coronary intervention. Those patients and 23 patients who did not attempt to 6-months follow up were excluded from the study. The remaining 304 patients comprise the study cohort. At 6-months follow-up, all patients were evaluated for symptoms by interview and

Table 2: Pre and post stent implantation echocardiographic parameters

Parameters	Baseline	Follow-up	P
Left atrium, cm	3.52±0.3	3.30±0.3	0.003
IVS, cm	0.97±0.1	0.95±0.1	NS
LV EDVI, ml/m ²	85.6±18.9	80.1±17.1	<0.0001
LV ESVI, ml/m ²	40.0±15.8	34.1±14.3	<0.0001
LV EF, %	53.2±11.9	57.0±11.1	<0.0001

IVS: interventricular septum, LV: left ventricular, EDVI: end-diastolic volume index, ESVI: end-systolic volume index, EF: ejection fraction.

examination and, echocardiography was repeated to assess recovery of LV function.

Statistics: The baseline characteristics of the population were analyzed using summary statistical measures. Categorical variables were presented numerically and continuous variables were presented as mean ± SD. The change in echocardiographic parameters before and 6 months after stent implantation was examined by paired Student t test, and the differences in LVEF changes between groups were examined by unpaired Student t test. The Chi square test was used for testing categorical variables. To identify "independent" prognostic factors of LVEF improvement, multivariate linear regression analyses were performed. All measures with a univariate p value <0.20 were considered for inclusion into multivariate analysis. A p value <0.05 was considered significant. All statistical analyses were performed using commercially available SPSS version 10.0.

RESULTS

Baseline characteristics: Baseline characteristics of the patients are summarized in in Table I. Most patients were men (87.9%) with a mean age of 60.4±10.8 years. Nineteen percent of the patients had diabetes mellitus (DM) and 44.0% of the patients was under antihypertensive treatment. About 55% of patients had a previous MI in the territory of the occluded artery. The mean duration of occlusion was 6.1±7.0 months. The occluded vessel was the left anterior descending artery in 40.8%, the left circumflex artery in 27.3% and the right coronary artery in 31.9%. In 77.3%, a single stent and in 22.7%, more than one stent was used per patient. The implanted stent type was BMS in 54.3% and DES in 45.7%.

Echocardiography: At 6-month follow-up we observed a significant improvement in LV function as shown by a decrease in LV EDVI and LV ESVI, and an increase in LVEF (p<0.0001) (Table II). Left atrial diameter was significantly decreased after 6-month

follow up (p<0.005).

Univariate analysis of change in LVEF in study subgroups:

Table III summarizes the univariate comparisons of change in LVEF between different study subgroups. After 6-month follow up a greater recovery of LVEF was detected in non-diabetic patients compared to those diabetic patients (p<0.04). Improvement in LVEF was observed both in patients with (p<0.05) and without hypertension (p<0.001). Although the treatment with angiotensin converting enzyme (ACE) inhibitors was associated with a greater magnitude of improvement in LVEF, the difference between patients with ACE inhibitors treatment and those without ACE inhibitors treatment did not reach statistical significance (P=0.06). A significant improvement in LVEF was observed from baseline to 6-month follow-up in both Canadian Cardiovascular Society (CCS) angina class 1, 2 and CCS class 3, 4 groups (p<0.001 and p<0.05 respectively). The intergroup differences in improvement of LVEF were not statistically significant. Regarding the baseline LVEF, patients with baseline LVEF <50% showed a significant improvement in LVEF while no significant changes were observed in patients with baseline LVEF >50% (p<0.0001 and p>0.05 respectively). After 6-month follow up a greater recovery of LVEF was detected in patients with occlusion duration <2 months than those with duration >2 months (p<0.01). No difference in the improvement of LVEF was noted between BMS or DES groups. However, the patients receiving multiple stents showed a greater improvement in LVEF than those patients with single stent implantation (p<0.05).

Multivariate predictors of improvement in LVEF: Multivariate regression analysis showed that baseline LVEF <50%, DM absence and occlusion duration <2 month to be independent predictors of improvement in LVEF (Table IV).

DISCUSSION

Successful PTCA of CTO and improvement in LV recovery after percutaneous revascularization has

Table 3: Comparison of baseline and six-month follow-up LV EF results according to variables

Variable	Baseline LV EF (%)	Follow-up LV EF (%)	p Value
Male gender			
Yes (n= 268)	53.1±12.2	56.9±12.2	0.926
No (n= 36)	53.8±11.7	57.7±11.0	
Diabetes mellitus			
Yes (n= 58)	51.5±12.0	53.8±11.8	0.040
No (n= 246)	53.6±12.1	57.7±11.7	
Treated hypertension			
Yes (n= 135)	50.9±12.5	53.9±12.2	0.119
No (n= 169)	55.0±11.1	59.5±10.8	
Previous myocardial infarction			
Yes (n= 168)	50.2±12.7	53.9±12.6	0.873
No (n= 136)	56.9±10.8	60.8±11.0	
Treatment of ACE inhibitors			
Yes (n = 114)	51.4±13.3	56.1±12.7	0.058
No (n = 190)	54.3±11.2	57.5±11.9	
Treatment of nitrate			
Yes (n = 182)	50.9±13.6	55.0±13.7	0.381
No (n = 122)	56.6±11.8	60.0±11.5	
Treatment of blockers			
Yes (n = 186)	53.8±11.6	57.9±11.3	0.342
No (n = 118)	52.2±12.3	55.6±12.6	
Treatment of calcium antagonist			
Yes (n = 92)	55.7±12.5	58.6±11.9	0.217
No (n = 212)	52.1±12.8	56.3±12.3	
Treatment of statin			
Yes (n = 142)	52.6±12.1	56.7±12.5	0.460
No (n = 162)	53.7±11.9	57.3±12.3	
Baseline angina class			
CCS class 1, 2 (n = 155)	54.0±12.2	58.4±11.6	0.156
CCS class 3, 4 (n = 149)	52.4±11.8	55.5±11.3	
Occlusion duration			
<2 month (n = 158)	52.1±12.1	57.2±11.0	0.004
>2 month (n = 146)	54.4±11.8	56.8±11.3	
Baseline LV EF (%)			
<50 (n = 141)	43.7±8.9	51.0±10.5	<0.0001
>50 (n = 163)	61.4±6.8	62.2±6.6	
Collateral circulation			
Yes (n = 186)	53.1±11.9	56.8±12.0	0.838
No (n = 118)	53.4±11.8	57.3±12.1	
Target-lesion coronary artery			
Left anterior descending (n = 121)	52.3±12.7	55.9±12.5	0.692
Left circumflex (n = 84)	54.2±11.2	58.2±11.0	
Right coronary (n = 99)	53.4±11.1	57.3±11.0	
Stents per patient			
1 (n = 235)	54.6±11.4	57.8±10.9	0.012
>2 (n = 69)	48.4±11.8	54.2±11.6	
Stent type			
Standard stent (n= 165)	53.4±11.7	56.7±11.3	0.197
Drug-eluting stent (n= 139)	52.9±11.9	57.4±10.6	

ACE: angiotensin-converting enzyme, CCS: Canadian cardiovascular society, LV EF: left ventricular ejection fraction

been demonstrated in several studies^{11,12,14}. PTCA, however, associated with a high rate of restenosis and late vessel occlusion. Recent studies have

shown that improvement in LV function has indeed observed only in patients without reocclusion^{11,12}. The stent implantation following percutaneous recanaliza-

Table 4: Multivariate predictors of improvement in LV EF

Predictors	Coefficient	p Value
Hypertension existence	-0.213	0.098
Diabetes mellitus existence	-0.245	0.027
ACE inhibitors use	0.274	0.122
Baseline angina class (CCS class 1, 2)	0.223	0.246
Occlusion duration (<2 month)	0.282	0.017
Baseline LV EF (< 50)	0.349	<0.0001
Multiple stent implantation	0.230	0.135
Drug-eluting stent implantation	0.280	0.180

ACE: angiotensin-converting enzyme, CCS: Canadian cardiovascular society, LV EF: left ventricular ejection fraction

tion of CTO decreases restenosis and reocclusion rates and therefore, increases the overall improvement in LV function^{8,9,17,18}. In accordance with previous reports we demonstrated both a reduction in LV volume and a significant improvement in LV function at follow-up after successful recanalization of CTO^{8,9,11}. We also confirmed that the most powerful determinant of improvement in LVEF at 6-month follow up was baseline LVEF <50%^{9,11}. Previous reports has been suggested that poor LV function in patients with coronary artery disease might be associated with perfusion defects^{9,11,20}. Such patients may have stunned or hibernating viable myocardium that recover after revascularization. Moreover, ventricles with low LVEF have much room to improve further after revascularization of a coronary occlusion.

In our study, multivariate analysis showed duration of occlusion to be independently predictive of improvement in LV function. There have been only two studies in literature assessing the relation between duration of coronary occlusion and improvement of LV function after percutaneous revascularization of CTO^{9,11}. Dzavik et al. demonstrated that duration of occlusion was independently predictive of improvement in LV function in patients with CTO submitted to percutaneous revascularization, as was the case in the present study⁹. In contrast to those results, however, Sirnes et al. did not found any relation between duration of coronary occlusion and LV recovery after percutaneous revascularization of CTO¹¹. Sirnes et al. conducted their study in a relatively small study population and 29 % of the patients underwent PTCA without stent implantation. This may be the reason for a diminished ability to detect a relationship between duration of coronary occlusion and recovery of LV function. Ours and Dzavik et al.'s findings indicate that dysfunctional but viable myocardium may persist several months after coronary artery occlusion, suggesting that reperfusion of

the target artery within 2 months may restore the persistent hibernating myocardium leading to recovery of LV function and preventing ventricular dilatation.

In the present study DM appears to be independently predictive of improvement in LV function. Our results are consistent with those of previous reports^{20,21}. Sugioka et al. indicated that CABG was inadequate for improving LV function in diabetic patients with severe ventricular dysfunction at the baseline²¹. Hu et al. reported that LVEF improvement was observed in non-diabetic patients but not in diabetic patients after percutaneous revascularization of left main coronary artery²⁰. It appears that DM itself may cause metabolic myocardial damage and microvascular functional abnormalities and thus may impair the recovery of LVEF after coronary revascularization. Moreover, higher restenosis rate after percutaneous stent implantation in diabetic patients compared with those non diabetic patients may also affect improvement of LV function after percutaneous revascularization²².

Regarding CSS class, Dzavik et al. reported that LVEF was improved more in patients with less severe angina⁹. However, Sirnes et al. reported that patients with angina pectoris CCS class 0-2 before PTCA had the same increase in LVEF as those with CCS class >2¹¹. The results of the present study are in accordance with the results of Sirnes et al. We found similar increase in LVEF in patient with angina pectoris class CCS class 1-2 and CCS class 3-4. It seems that CSS class is not a predictive parameter in LV EF recovery after coronary revascularization.

There are conflicting reports on the influence of collaterals on recovery of LV EF in coronary occlusions. Some authors have reported that the presence of well-developed collaterals was an important factor in determining LV function in patients with coronary occlusions^{23,24}. In contrast, other authors have reported no effect of collaterals on baseline function

or improvement of LVEF after reperfusion²⁵. In our study, we found that collateral circulation had no influence on LV recovery after revascularization in patients with coronary occlusions. It appears that the presence of coronary collaterals is not mandatory for the functional recovery in patients with CTO.

Many authors have reported the MI to be a factor affecting LVEF after coronary revascularization^{5,26}. In those studies it was stated that patients with a history of MI would have less viable ischemic myocardium and, thus it would be less likely to benefit from coronary revascularization. On the other hand, some other authors have reported that there was no significant difference in recovery of LVEF after revascularization of coronary occlusion between patients with old MI and those without previous MI^{9,11}. In accordance with later reports, we did not found significant difference in recovery of LVEF between patients with history of MI >2 months and those without history of MI. The reason for these discrepancies may be explain with heterogeneous of the study groups.

DES implantation, which enhances long-term patency of the occluded vessel compared with BMS implantation, might be advantageous²⁷. There is only one study in literature comparing improvement of LV function after DES and BMS implantation in patients with CTO. In this study, Nakamura et al. have reported that improvement in LV function after DES stent implantation was better than BMS implantation⁷. In our study, we found a higher increment of EF in patients revascularized with DES stents compared with BMS stents, however, the difference that did not reach statistical significance. This may have been in part because we performed follow-up echocardiography at 6-month and most of restenosis were detected at an earlier stage and, therefore, only a small percentage of restenotic lesions were found to be severely narrowed which may influence LV function.

It is well known that ACE inhibitor improves LVEF in patients with LV dysfunction^{28,29}. In the present study drug treatment with ACE inhibitor did not enhance recovery of LV function in multivariate analysis. The reason for this contrary may be due to the patients in our study had less LV systolic dysfunction as compared with the patients in the large ACE inhibitor trials^{28,29}. Moreover, the effect of ACE inhibitor on improvement of LV function seems most striking in the early months after initiation of treat-

ment. Most of the patients in our study had started ACE inhibitors months before percutaneous coronary stent implantation.

LIMITATIONS

There are several limitations of this study. First, we did not have a control group with no intervention; thus no treatment effect on LV function can be ascribed directly to the revascularization procedure. Second, coronary angiography was not repeated at 6-months follow up; thus, restenosis could not be excluded. Third, all patients in this study had angina pectoris or a positive exercise stress test. However, many patients with CTO have no myocardial ischaemia related to the occluded artery. Thus, the possible benefit in LV function after stent implantation of CTO in ischaemic patients demonstrated in this study cannot be generalized to patients without ischaemia.

CONCLUSION

Stent implantation of CTO improves LVEF and decreases LV volumes during 6 months follow-up after coronary stent implantation. Improvement in LV function is most pronounced in patients with low baseline LVEF and those with coronary occlusion within 2 months. The presence of diabetes mellitus impairs recovery of LVEF after coronary revascularization. However, CSS class, presence of coronary collaterals or history of MI >2 months have no effect on improvement of LV function.

REFERENCES

1. Hammermeister KE, DeRouen TA, Dodge HT. Variables predictive of survival in patients with coronary disease. Selection by univariate and multivariate analyses from the clinical, electrocardiographic, exercise, arteriographic, and quantitative angiographic evaluations. *Circulation* 1979; 59:421-30.
2. White HD, Norris RM, Brown MA, Brandt PW, Whitlock RM, Wild CJ. Left ventricular end-systolic volume as the major determinant of survival after recovery from myocardial infarction. *Circulation* 1987; 76:44 -51.
3. Georgievska-Ismail L, Vavlukis M, Borozanov V. Predictive value of left ventricular function on prognosis in patients undergoing coronary artery bypass surgery. *Prilozi* 2005; 26:103-19.

4. Bax M, de Winter RJ, Schotborgh CE, Koch KT, Meuwissen M, Voskuil M, et al. Short- and long-term recovery of left ventricular function predicted at the time of primary percutaneous coronary intervention in anterior myocardial infarction. *J Am Coll Cardiol* 2004; 43:534-41.
5. Halkin A, Stone GW, Dixon SR, Grines CL, Tchong JE, Cox DA, et al. Impact and determinants of left ventricular function in patients undergoing primary percutaneous coronary intervention in acute myocardial infarction. *Am J Cardiol* 2005; 96:325-31.
6. Miketic S, Carlsson J, Tebbe U. Improvement of global and regional left ventricular function by percutaneous transluminal coronary angioplasty after myocardial infarction. *J Am Coll Cardiol* 1995; 25:843-47.
7. Nakamura S, Muthusamy TS, Bae JH, Cahyadi YH, Udayachalerm W, Tresukosol D. Impact of sirolimus-eluting stent on the outcome of patients with chronic total occlusions. *Am J Cardiol* 2005; 95:161-66.
8. Van Belle E, Blouard P, McFadden EP, Lablanche JM, Bauters C, Bertrand ME. Effects of stenting of recent or chronic coronary occlusions on late vessel patency and left ventricular function. *Am J Cardiol* 1997; 80:1150-54.
9. Dzavik V, Carere RG, Mancini GB, Cohen EA, Catellier D, Anderson TE, et al.; Total Occlusion Study of Canada Investigators. Predictors of improvement in left ventricular function after percutaneous revascularization of occluded coronary arteries: a report from the Total Occlusion Study of Canada (TOSCA). *Am Heart J* 2001; 142:301-308.
10. Baks T, van Geuns RJ, Duncker DJ, Cademartiri F, Mollet NR, Krestin GP, et al. Prediction of left ventricular function after drug-eluting stent implantation for chronic total coronary occlusions. *J Am Coll Cardiol* 2006; 47:721-25.
11. Sirnes PA, Myreng Y, Molstad P, Bonarjee V, Golf S. Improvement in left ventricular ejection fraction and wall motion after successful recanalization of chronic coronary occlusions. *Eur Heart J* 1998; 19:273-81.
12. Piscione F, Galasso G, De Luca G, Marrazzo G, Sarno G, Viola O, et al. Late reopening of an occluded infarct related artery improves left ventricular function and long term clinical outcome. *Heart* 2005; 91:646-51.
13. Engelstein E, Terres W, Hofmann D, Hansen L, Hamm CW. Improved global and regional left ventricular function after angioplasty for chronic coronary occlusion. *Clin Investig* 1994; 72:442-47.
14. Puma JA, Sketch MH Jr, Tchong JE, Harrington RA, Phillips HR, Stack RS, et al. Percutaneous revascularization of chronic coronary occlusions: an overview. *J Am Coll Cardiol* 1995; 26:1-11.
15. Berger PB, Holmes DR Jr, Ohman EM, O'Hanesian MA, Murphy JG, Schwartz RS, et al. Restenosis, reocclusion and adverse cardiovascular events after successful balloon angioplasty of occluded versus nonoccluded coronary arteries. Results from the Multicenter American Research Trial With Cilazapril After Angioplasty to Prevent Transluminal Coronary Obstruction and Restenosis (MARCATOR). *J Am Coll Cardiol* 1996; 27:1-7.
16. Violaris AG, Melkert R, Serruys PW. Long-term luminal renarrowing after successful elective coronary angioplasty of total occlusions. A quantitative angiographic analysis. *Circulation* 1995; 91:2140-50.
17. Sirnes PA, Golf S, Myreng Y, Molstad P, Emanuelsson H, Albertsson P, et al. Stenting in Chronic Coronary Occlusion (SICCO): a randomized, controlled trial of adding stent implantation after successful angioplasty. *J Am Coll Cardiol* 1996; 28:1444-51.
18. Ashby DT, Dangas G, Mehran R, Leon MB. Coronary artery stenting. *Catheter Cardiovasc Interv* 2002; 56:83-102.
19. American College of Cardiology/American Heart Association task force on practice guidelines (Committee to revise the 1993 guidelines for percutaneous transluminal coronary angioplasty); Society for Cardiac Angiography and Interventions. ACC/AHA guidelines for percutaneous coronary intervention (revision of the 1993 PTCA guidelines)-executive summary: a report of the American College of Cardiology/American Heart Association task force on practice guidelines (Committee to revise the 1993 guidelines for percutaneous transluminal coronary angioplasty) endorsed by the Society for Cardiac Angiography and Interventions. *Circulation* 2001; 103:3019-41.

20. Hu FB, Tamai H, Kosuga K, Kyo E, Hata T, Okada M, Nakamura T, Fujita S, Tsuji T, Takeda S, Motohara S, Uehata H. Predictors of improvement in left ventricular function after initially successful angioplasty of unprotected left main coronary artery stenoses. *Int J Cardiovasc Intervent* 2004; 6:119-27.
21. Sugioka J, Ozawa S, Inagaki M, Fukuzawa S, Daimon M, Kushida S, et al. Influence of diabetes mellitus on left ventricular function in patients undergoing coronary artery bypass grafting. *J Cardiol* 2000; 36:9-16.
22. Weisz G, Leon MB, Holmes DR Jr, Kereiakes DJ, Clark MR, Cohen BM, et al. Two-year outcomes after sirolimus-eluting stent implantation: results from the Sirolimus-Eluting Stent in de Novo Native Coronary Lesions (SIRIUS) trial. *J Am Coll Cardiol* 2006; 47:1350-55.
23. Fukui M, Li M, Nakakoji T, Kawakatsu M, Nariyama J, Yokota N, et al. Angiographically demonstrated coronary collaterals predict residual viable myocardium in patients with chronic myocardial infarction: a regional metabolic study. *J Cardiol* 2000; 35:103-11.
24. Juilliere Y, Marie PY, Danchin N, Karcher G, Bertrand A, Cherrier F. Evolution of myocardial ischemia and left ventricular function in patients with angina pectoris without myocardial infarction and total occlusion of the left anterior descending coronary artery and collaterals from other coronary arteries. *Am J Cardiol* 1991; 68:7-12.
25. Di Carli M, Sherman T, Khanna S, Davidson M, Rokhsar S, Hawkins R, et al. Myocardial viability in asynergic regions subtended by occluded coronary arteries: relation to the status of collateral flow in patients with chronic coronary artery disease. *J Am Coll Cardiol* 1994; 23:860-68.
26. Chung CM, Nakamura S, Tanaka K, Tanigawa J, Kitano K, Akiyama T, et al. Effect of recanalization of chronic total occlusions on global and regional left ventricular function in patients with or without previous myocardial infarction. *Catheter Cardiovasc Interv* 2003; 60:368-74.
27. Migliorini A, Moschi G, Vergara R, Parodi G, Carrabba N, Antoniucci D. Drug-eluting stent-supported percutaneous coronary intervention for chronic total coronary occlusion. *Catheter Cardiovasc Interv* 2006; 67:344-48.
28. Kober L, Torp-Pedersen C, Carlsen JE, Bagger H, Eliassen P, Lyngborg K, et al. A clinical trial of the angiotensin-converting-enzyme inhibitor trandolapril in patients with left ventricular dysfunction after myocardial infarction. Trandolapril Cardiac Evaluation (TRACE) Study Group. *N Engl J Med* 1995; 333:1670-76.
29. Flather MD, Yusuf S, Kober L, Pfeffer M, Hall A, Murray G, et al. Long-term ACE-inhibitor therapy in patients with heart failure or left-ventricular dysfunction: a systematic overview of data from individual patients. ACE-Inhibitor Myocardial Infarction Collaborative Group. *Lancet* 2000; 355:1575-81.